


# SURFICIAL GEOLOGIC MAP OF THE SOUTHWICK QUADRANGLE, CLEARWATER AND NEZ PERCE COUNTIES, IDAHO

Kurt L. Othberg, Daniel W. Weisz, and Roy M. Breckenridge  
2002

**Disclaimer:** This Digital Web Map is an informal report and may be revised and formally published at a later time. Its content and format may not conform to agency standards.



## DESCRIPTION OF MAP UNITS

### INTRODUCTION

The surficial geologic map of the Southwick quadrangle identifies earth materials on the surface and in the shallow subsurface. It is intended for those interested in the area's natural resources, urban and rural growth, and private and public land development. The information relates to assessing diverse conditions and activities, such as slope stability, construction design, sewage drainage, solid waste disposal, and ground-water use and recharge.

The geology was intensively investigated during a one-year period. Natural and artificial exposures of the geology were examined and selectively collected. In addition to field investigations, aerial photographs were studied to aid in identifying boundaries between map units through photogeologic mapping of landforms. In most areas map-unit boundaries (contacts) are approximate and were drawn by outlining well-defined landforms. It is rare that contacts between two units can be seen in the field without excavation operations which are beyond the purpose and scope of this map. The contacts are inferred where landforms are poorly defined and where lithologic characteristics grade from one map unit into another. The precision of a contact with respect to actual topography also depends on the accuracy and scale of the topographic base. Details depicted at this scale, therefore, provide an overview of the area's geology. Further intensive analyses at specific locations should be arranged through independent geotechnical specialists.

Teakam Butte and the canyons of the Clearwater River and two tributaries, Bedrock and Louse creeks, are the prominent features on the map. The Clearwater River drains the northern Rocky Mountains and flows through Teakam Butte is part. The river's canyon exposes the plateau's Miocene basalt flows of the Columbia River Basalt Group and the underlying pre-Miocene granitic and metamorphic rocks that compose the nearby Northern Rocky Mountains. During the Miocene, lava flows of the Columbia River Basalt Group filled ancestral stream valleys eroded into the basement rocks. The flows created volcanic embayments that now form the eastern edge of the Columbia River Plateau where the relatively flat region trends the mountains. Sediments of the Latah Formation are interbedded with the basalt flows, and landslide deposits occur where major sedimentary interbeds are exposed along the valley sides. Pleistocene loess forms a thin discontinuous mantle on deeply weathered surfaces of the basalt plateau and mountain foothills. In the late Pleistocene, multiple Lake Missoula Floods inundated the Clearwater River valley, locally depositing silt, sand, and ice-raffed pebbles and cobbles in the lower elevations of the canyon.

The bedrock geology of this area is mapped by Lewis and others (2001) and shows details of the basement rocks and the Miocene basalt flows and sediments. The bedrock map's cross sections are especially useful for interpreting subsurface conditions suitable for siting water wells and assessing the extent and limits of ground water.

### SURFICIAL DEPOSITS

**Qam** Alluvium of mainstreams (Holocene)—Channel and flood-plain deposits of the Clearwater River that are actively being formed on a seasonal or annual basis. Two grain-size suites are typically present: Well-sorted and rounded sandy gravel of river bars and islands, and coarse sand forming thin shoreline deposits. The gravel includes clasts of basaltic, granitic, and metamorphic rocks. These deposits are called riverwash-argents and Aquic Xenofluids in the soil surveys (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999; Glenn Hoffman, written comm., 2001).

**Qoam** Older alluvium of mainstreams (Holocene)—Fine to coarse-grained bedded sand and silty sand overlying river channel gravel. These alluvial deposits form one or more levels of old point bars and flood plains of the Clearwater River which are younger than the Lake Missoula Floods backwater deposits, but older than alluvium of the present river. Surface heights above present mean water level range from 9' to 45' feet. Relative heights and soil characteristics suggest a late Holocene age, and the lower of these surfaces may have been inundated by the highest seasonal flood waters before the stream flows were controlled by Dworshak dam. The sand overlying channel gravel is several feet thick. Soils developed in older mainstream alluvium include the Uhlug series (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999).

**Qas** Alluvium of sidestreams (Holocene)—Channel and flood-plain deposits of Louse Creek. Primarily coarse channel gravels deposited during high-energy stream flows. Subrounded to rounded pebbles, cobbles, and boulders of basalt in a sand matrix. Moderately stratified and sorted. Includes intercalated colluvium and debris-flow deposits from steep side slopes.

**Qac** Alluvium and colluvium (Holocene)—Stream, slope-wash, and gravity deposits. Predominantly beds of silt, clay, and sand derived from erosion of adjacent units. Stream deposits typically are thin and intertongued with laterally thickening deposits of slope wash and colluvium derived from local debris deposits and weathered basalt. Soils developed in these deposits include the Itzee, Lovell, Latacho, Teneb, and Wilkins series (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999; Glenn Hoffman, written comm., 2001).

**Qad** Alluvial fan and debris-flow deposits (Holocene and Pleistocene)—Primarily crudely bedded, poorly sorted brown muddy gravel shed from canyon slopes of basalt colluvium. Gravel is composed of subangular and angular pebbles, cobbles, and boulders of basalt in a matrix of granules, sand, silt, and clay. May include beds of silt and sand derived from reworked loess, Mazama ash, and Lake Missoula Flood backwater deposits. Thickness varies but typically ranges from 6-50 feet. Fans composed of alluvium and debris flows occur in canyon bottoms below steep debris-flow chutes (see Symbols).

**Qle** Landslide deposits (Holocene and Pleistocene)—Poorly sorted and poorly stratified angular basalt cobbles and boulders mixed with silt and clay. Landslide deposits include debris slides as well as blocks of basalt, sedimentary interbeds, and pre-Miocene rocks that have been rotated and moved laterally. Debris slides mainly composed of unstratified, unsorted gravel rubble in a clayey matrix. In addition to the landslide deposit, the unit may include the landslide scarp and the headwall area adjacent to and below the landslide scarp from which material broke away (see Symbols). The headwall area may include talus formed after landslide movement. Location of landslide deposits in canyons is controlled by the presence of sedimentary interbeds, the hydrologic regime, and the occurrence of basalt overlying clay-rich weathered basement rocks. The largest landslides occur where canyon-cutting has exposed landslide-prone sediments to steep topography. Slope failures have occurred where the fine-grained sedimentary interbeds and weathered basement rocks are saturated by ground water moving toward the valleys. This relationship is so prevalent that the major sedimentary interbeds may be traced by locating landslide deposits along the valley sides. The landslides range in age from ancient, relatively stable features to those that have been active within the past few years. The factors that cause landslides have been prevalent in the region for thousands of years. The frequency of landsliding may have been greater in the Pleistocene. Today, initiation and reactivation of landslides is closely tied to unusual climatic events and land-use changes. Even small landslide activity on the upper parts of canyon slopes can transform into high-energy debris flows that endanger roads, buildings, and people below (see Debris-flow chute under Symbols). Landslide debris is highly unstable when modified through natural variations in precipitation, artificial cuts, fills, and changes to surface drainage and ground water.

**Qcb** Colluvium from basalt (Holocene and Pleistocene)—Primarily poorly sorted brown muddy gravel composed of angular and subangular pebbles, cobbles, and boulders of basalt in a matrix of silt and clay. Emplaced by gravity movements on steep-sided canyons and gullies cut into Columbia River basalt. Includes outcrops of basalt that are common on steep, dry, southerly aspects where colluvium is thinner and the more erosion-resistant basalt flows form laterally traceable ledges. More gently sloping areas are mantled with thin loess (typically 1-5 feet thick), especially near boundaries with loess mantling Tertiary residuum (QTbr). Distribution and thickness of colluvium is dependent on slope aspect, upper and lower slope position, basalt and sediment stratigraphy, and association with landslides. Colluvium is thin and associated with many basalt outcrops on dry, southerly facing slopes, and may exhibit patterned-ground features (see Symbols). Colluvium is thicker on north- and east-facing slopes, and is associated with landslides (Qb) and debris-flow chutes (see Symbols), especially where more moisture is retained and where sedimentary interbeds are present. Areas of thicker colluvium have fewer outcrops of basalt, and the surface may have a patterned ground of recent-shaped beds of colluvium, probably relict of Pleistocene solifluction. Unit includes landslides too small to map separately, and talus below cliffs and ledges of basalt. Colluvium typically increases in thickness toward the base of slopes where it intertongues with alluvium in valley bottoms. May include soil of valley-bottom sediment where streams have little discharge or are ephemeral. Soils developed in basalt colluvium include the Agatha, Gwin, Hooverton, Jacket, Ketenback, Keaterville, Klicson, Meland, Upton, and Wellbech series (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999; Glenn Hoffman, written comm., 2001).

**Qog** Colluvium from granitic and metamorphic rocks (Holocene and Pleistocene)—Primarily poorly sorted muddy gravel composed of angular and subangular pebbles, cobbles, and boulders in a matrix of silt, sand, and clay. Emplaced by gravity movements in canyons where there are outcrops of pre-Miocene granitic and gneissic rocks. Includes local debris-flow deposits and isolated rock outcrops. May include colluvium and debris-flow deposits from the upslope basalt section, and areas of thin loess (typically less than 5 feet). Grades laterally, as slope gradients decrease, into areas where thin loess and clayey saprolite mantle bedrock. Distribution and thickness of colluvium depend on slope, aspect, upper and lower slope position, and association with landslide deposits. Colluvium typically increases in thickness toward the base of slopes and may intertongue with alluvium in valley bottoms. Soils developed in these deposits include the Aldermard, Rudo, Teascreek, and Whiskeycreek series (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999; Glenn Hoffman, written comm., 2001).

**Qag** Lake Missoula Floods backwater deposits (Pleistocene)—Rhythmites deposited when Lake Missoula Floods backwaters inundated the Clearwater River valley. Primarily alternating thin beds of gray sand and pale brown silt. Cross-bedded, dark-gray, basalt-rich granule gravel and coarse sand may be present at the base. Includes cut and fill structures and sandy elastic dikes. Similar depositional environment, sedimentology, and age as Lake Missoula Floods rhythmites of eastern Washington (Smith, 1993; Waitt, 1980, 1985). Commonly reworked into sandy, silty colluvium. Found locally up to 1,200 feet in elevation, the approximate maximum flood level. Mapped as a pattern where sandy, silty rhythmites mantle deposits of debris flows and alluvial fans (Qad) and Pleistocene alluvial gravel (Qag). Downstream at Lewiston in the Snake River valley, Lake Missoula Floods backwater deposits overlie Bonneville Flood gravel. In the Clearwater River drainage, Bonneville Flood deposits have not been recognized. Lake Missoula Floods temporarily reversed the course of the Clearwater River within the area of backwater inundation (see Flow direction in Symbols). Soils developed in Lake Missoula Flood deposits include the Uhlug series (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999).

**Qag** Alluvial gravel (Pleistocene)—Well-rounded pebble and cobble gravel of remnant point bars that range 52 to 78 feet above the Clearwater River. Gravel poorly exposed owing to mantle of Lake Missoula Floods backwater sediments (Qm). Intertongues with colluvium and debris-flow deposits at toe of canyon slope. The gravel was deposited by the ancestral Clearwater River prior to the latest Lake Missoula Floods and may have formed during periodic greater discharges of the river during the Wisconsin glaciation. Soils mapped in areas of Pleistocene alluvial gravel include the Uhlug series (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999).

**QTcr** Colluvium and residuum from metamorphic rocks (Quaternary and Tertiary)—Primarily poorly sorted muddy gravel composed of angular and subangular pebbles, cobbles, and boulders in a matrix of sand, silt, and clay. Grades laterally, as slope gradients decrease, into areas where matrix predominates and areas of saprolite associated with Tertiary sediments. Parent rocks include schist, gneiss, and quartzite. In canyons includes isolated outcrops of metamorphic rocks and debris-flow deposits from the overlying basalt section. Distribution and thickness depend on degree of weathering of parent rock, slope, aspect, and upper and lower slope position. Rock outcrops and gravelly colluvium are more common on steep slopes and at higher elevations, and rare at lower elevations and gentle slopes. At lower elevations a mantle of thin loess is common (typically less than 5 feet). Soils developed in these deposits include the Dworshak, Hugas, Kruse, McCrosket, and Teakcam series (Glenn Hoffman, written comm., 2001).

**QTbr** Loess and basalt residuum (Quaternary and Tertiary)—Thin Quaternary loess mantling Tertiary residuum on remnant surfaces of the basalt plateau. Loess 1-6 feet thick mantles basalt which is spheroidally weathered and commonly grades upward into a thoroughly decomposed zone of clayey saprolite. Most weathered saprolite have indicated cores of basalt which grade outward into yellowish and reddish sand, silt, and clay. Original fracture patterns in the basalt can be seen as veins of secondary accumulation of clay, iron oxides, and, locally, opal. The saprolite varies from yellowish brown to dark red in color and predominantly consists of silty clay and clay. The saprolite mineralogy consists predominantly kaolinite clay and oxides of iron, especially goethite and hematite. The basalt residuum is laterally discontinuous, probably as a result of erosion of the Miocene land surface, so that near drainages and canyon rims fresh basalt is often near or at the present surface. Where thickest, the clayey saprolite is as much as 20 feet thick and spheroids may not be seen in road cuts or near the land surface. Elsewhere, the saprolite is thinner, and spheroidally weathered basalt is seen in road cuts and lag boulders of weathered spheroids are encountered in fields. The weathering of the basalt probably can be attributed to the eastward increase in precipitation and to the Miocene age of this remnant basalt surface. Includes gravelly basalt colluvium on local steeper slopes where stream incision has occurred and local deposits of thin alluvium too small in area to show at this scale. Soils in this unit include the Cavendish, Joel, Larkin, Southwick, and Tany series (U.S. Department of Agriculture, Natural Resources Conservation Service, 1999; Glenn Hoffman, written comm., 2001).

**QTlr** Loess, sediment, and residuum (Quaternary and Tertiary)—Thin Quaternary loess mantling Tertiary transported and residual deposits of sand, silt, and clay. Rare to common granules, pebbles, and cobbles may occur in thin beds or as isolated clasts. Within the unit, a gradation exists between (1) sediment transported from foothill and mountain sources to depositional sites adjacent to and on top of basalt along the margins of the basalt plateau, and (2) residual deposits formed in place by weathering of pre-Miocene rocks (QTer). In many places the sediments and residuum have been weathered to a clay-rich saprolite by the warm- and wet-climate of the late Tertiary. The presence of quartz grains distinguishes saprolite formed in sediments from basalt saprolite. Includes local deposits of thin alluvium too small to show at this scale. Soils in this unit include the Carlinton, Joel, and Tany series (Glenn Hoffman, written comm., 2001).


### REFERENCES

- Lewis, R.S., J.H. Bush, R.F. Burnester, J.D. Kaufman, D.L. Garwood, P.E. Meyers, K.L. Othberg, and W.C. McClelland, in preparation. Geologic map of the Potlatch 30' x 60' quadrangle, Idaho: Idaho Geological Survey Special Report, 1:100,000 scale.
- Smith, C.A., 1993. Missoula flood dynamics and magnitudes inferred from sedimentology of slackwater deposits on the Columbia Plateau, Washington: Geological Society of America Bulletin, v. 105, p. 77-100.
- U.S. Department of Agriculture, Natural Resources Conservation Service, 1999. Soil survey geographic (SSURGO) database for Clearwater and Nez Perce counties, USDA-NRCS Soil Survey Division, National SSURGO Database Data Access, ID611.
- Waitt, R.B., Jr., 1985. Case for periodic, colossal jökulhaups from Pleistocene glacial Lake Missoula: Geological Society of America Bulletin, v. 95, p. 1271-1286.
- Waitt, R.B., Jr., 1980. About 40 last-glacial Lake Missoula jökulhaups from southern Washington: Journal of Geology, v. 88, p. 653-679.

### ACKNOWLEDGEMENT

Glenn Hoffman (Soil Survey Party Leader, U.S. Department of Agriculture, Natural Resources Conservation Service, Orofino, Idaho) provided preliminary soil survey data for Clearwater County.

Base map from USGS digital raster graphic, 1990.  
Topography by photogrammetric methods from aerial photographs taken 1957. Field checked 1967. Revised from aerial photographs taken 1987.  
Polyconic projection, 1927 North American Datum.  
10,000-foot grid ticks based on Idaho coordinate system, west zone.  
100-meter Universal Transverse Mercator grid ticks, zone 11.  
National geodetic vertical datum of 1929.



Field work conducted 2001.  
This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program.  
Digital cartography by B. Benjamin E. Studer and Loudon R. Stanford at the Idaho Geological Survey's Digital Mapping Lab.  
Map version 7-8-2003.

### SYMBOLS

- Contact: Line showing the approximate boundary between one map unit and another. The apparent ground width of the line representing the contact is about 80 feet at this scale (1:24,000).
- Landslide scarp: Ticks show top of scarp.
- Debris-flow chute in canyons. Thin and discontinuous alluvial fan and debris flow deposits (Qad) may be present, but are not mappable at this scale. High-energy, short duration floods and debris flows may occur in these chutes in response to severe climatic conditions, such as thunderstorms and rain-on-snow events. These events are historically infrequent, dependent on weather, with a recurrence cycle on the order of years to decades. Debris flows can also be triggered by landslides. The most prominent debris-flow chutes are shown on the map, but any steep-gradient valley sides and canyon bottoms have the potential for these catastrophic events.
- Flow direction of Missoula Floods backwater inundation.
- Patterned ground associated with the weathered, differentially eroded surface of basalt. Pattern consists of regularly spaced, subround fracture system in basalt with silty mounds between fractures. Silty mounds give way to fractured basalt downslope, but thicken upslope where they gradually obscure the fracture pattern and merge with loess deposits or weathered basalt. Probably formed by stripping of loess from the basalt surface through periglacial processes. Original patterned ground features destroyed by field plowing in many locations.

### CORRELATION OF MAP UNITS

