


GEOLOGIC MAP OF THE FALLS CITY QUADRANGLE, JEROME COUNTY, IDAHO

John D. Kauffman and Kurt L. Othberg
2005

Disclaimer: This Digital Web Map is an informal report and may be revised and formally published at a later time. Its content and format may not conform to agency standards.

CORRELATION OF MAP UNITS



Sources of age determination:
* ⁴⁰Ar/³⁹Ar weighted mean age, Esser (2005)
† ⁴⁰Ar/³⁹Ar weighted mean av. stage plateau age, Tauxe and others (2004)

INTRODUCTION

The geologic map of the Falls City quadrangle identifies both the bedrock and surficial geologic units. It shows the geographic distribution of rock types at the surface and in the shallow subsurface. The Falls City quadrangle lies near the center of the Snake River Plain, a large, arcuate, lava-filled depression crossing southern Idaho. Pleistocene basalt flows from shield volcanoes, such as Flat Top Butte in this quadrangle, form the land surface. The older basalt flows are mantled with alluvium and wind-blown sand and silt which form the soils that are cultivated. Approximately 14,500 years ago the Bonneville Flood filled and overtopped the Snake River Canyon (O'Connor, 1993). Across the south part of the quadrangle, the Bonneville Flood stripped soils from the basalts of Rocky and Wilson buttes and locally deposited flood gravels. The geologic units in the area control soil development, groundwater movement and recharge, and geotechnical factors important in construction design and waste management. Land uses in the area include irrigated agriculture, rural and urban residential development, industrial and commercial enterprises, and dairy farms with confined animal feeding operations. The Snake River Plain aquifer underlies the area and discharges to the west of the Falls City quadrangle as springs in the Snake River Canyon.

Earlier geologic mapping by Malde and others (1963) was reviewed, and field checking of their map was combined with new field investigations in 2003-2004 of both bedrock and surficial geology. Exposures of the geology were examined and selectively sampled. Aerial photographs were studied to aid in identifying boundaries between map units through photogeologic mapping of landforms. The information depicted at this scale furnishes a useful overview of the area's geology but is not a substitute for site-specific evaluations.


DESCRIPTION OF MAP UNITS

- Qas Alluvium of side-streams (Holocene and Pleistocene)**—Silt and sand flood-plain and sheet-wash deposits in the drainage systems formed between older and younger basalt units.
- Bonneville Flood**
 - Qabf Scabland of flood pathways (Pleistocene)**—Flood-scoured basalt surface. Loess stripped, basin and butte topography is common. Unit adapted from Scott (1982) and O'Connor (1993). Original basalt morphology stripped of pre-flood loess and soils, in contrast with area north of maximum flood extent but in same basalt unit. Includes patchy sheets and bars of thin sand and gravel that are not mapped, but have been locally used for gravel (see Symbols).
 - Qabf Fine-grained deposits in slack-water basins (Pleistocene)**—Sand and silt deposited in basins of basalt surface that were protected from high-energy water flow.
- EOLIAN DEPOSITS**
 - Qwb Dune sand (Holocene)**—Thin, stratified fine sand of stabilized wind dunes. Shown only where identified on aerial photographs.

BASALT UNITS


- Qwb Basalt of Wilson Butte (Pleistocene)**—Dark gray to black, fine-grained basalt with common to abundant plagioclase phenocrysts 1-3 mm in length and fairly common olivine grains up to 1 mm in diameter, and some plagioclase-olivine intergrowths. Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Source is Wilson Butte northeast of the Falls City quadrangle. Land surface is rough from common pressure ridges and has little or no drainage development. North of the maximum extent of the Bonneville Flood (see Symbols), thin loess covers the surface of the unit except for tops of pressure ridges, and soil caliche is present but generally thin and weakly developed (Baldwin, 1925; Ames, 2003). Some of the land is cultivatable; most of the area is sagebrush and grasses. Gruhn (1961) reports a radiocarbon date of 15,000 years on tumbled camel bones found inside a lava tube from Wilson Butte, constraining the eruption of the lava to before that time (Matthews, 2000).
- Qrb Basalt of Rocky Butte (Pleistocene)**—Fine-grained, dark gray to black, glassy basalt with common to abundant olivine grains 0.5-1 mm and clusters 1 to 3 mm in diameter. Common to abundant small plagioclase laths to about 1 mm in length. Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Erupted from a shield volcano located 22 miles northeast of the city of Twin Falls in the Eden NE topographic quadrangle, which shows a permanent horizontal control mark labeled "Rocky" at 4526 feet on the south rim of the vent (Sec. 14, T. 8 S., R. 20 E.). Equivalent to Sand Springs Basalt of Malde and Powers (1962), Malde and others (1963), Covington (1976), and Covington and Weaver (1990). Covington and Weaver (1990) identified the source as "Butte 4526". In the Falls City quadrangle, all of unit was scoured by Bonneville Flood (Qabf) and is mostly outcrop. Little of the land is cultivatable. Tauxe and others (2004) report an ⁴⁰Ar/³⁹Ar weighted mean plateau age of 0.095 Ma for "Sand Springs" basalt. Their sample location is on the north rim of the Snake River canyon near Shoshone Falls, 1.5 miles south of the Falls City quadrangle. The sample is from the unit we map as basalt of Rocky Butte.
- Qftb Basalt of Flat Top Butte (Pleistocene)**—Fine-grained, medium gray basalt with scattered to very abundant plagioclase-olivine intergrowths 4-7 mm, and olivine grains and clots 1-4 mm. Flows typically vesicular near the top and more dense in the center, but diktytaxitic throughout with abundant fine-grained plagioclase laths. Carbonate coatings and fillings common in voids but not pervasive. Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Erupted from the Flat Top Butte shield volcano. Equivalent to Thousand Springs Basalt of Malde and Powers (1962), Malde and others (1963), and Gillerman and Schiappa (1994, 2001). Tauxe and others (2004) report an ⁴⁰Ar/³⁹Ar weighted mean plateau age of 0.335 Ma for this unit (their sample s09, Thousand Springs Basalts). An ⁴⁰Ar/³⁹Ar weighted mean age of 0.33±0.8 Ma was obtained on our sample 02P002B (Esser, 2005). Topography contrasts with surface of the younger basalt flows. Almost no basalt pressure ridges rise above a nearly complete mantle of loess. Surface drainage is moderately developed. Thickness of mantle ranges 3-25 feet; commonly 3-12 feet thick. Soil caliche (duripan) is typically well developed within the soil profile (Ames, 2003) and at the soil-basalt contact, but the thickness of caliche is varies considerably. Most of the land is cultivatable.

SYMBOLS




REFERENCES

Ames, Dal, 2003. Soil survey of Jerome County and Part of Twin Falls County, Idaho: U.S. Department of Agriculture, Natural Resources Conservation Service, 391 pages, 67 sheets.
Baldwin, Mark, 1925. Soil Survey of the Twin Falls area, Idaho: U.S. Department of Agriculture, Bureau of Soils, Advance Sheets—Field Operations of the Bureau of Soils, 1921, p. 1367-1394, 1 plate.
Covington, H.R., 1976. Geologic map of the Snake River Canyon near Twin Falls, Idaho: U.S. Geological Survey Miscellaneous Field Studies Map MF-809, 2 sheets, scale 1:24,000.
Covington, H.R. and J.N. Weaver, 1990. Geologic map and profiles of the north wall of the Snake River Canyon, Jerome, Filer, Twin Falls, and Kimberly quadrangles, Idaho: U.S. Geological Survey Miscellaneous Investigations Series, Map I-1947-D, scale 1:24,000.
Esser, Richard P., 2005. ⁴⁰Ar/³⁹Ar geochronology results from volcanic rocks from Idaho: New Mexico Geochronology Research Laboratory Internal Report NMGR-IR-431, 10 p.
Gruhn, Ruth, 1961. Archaeology of Wilson Butte Cave, Idaho Museum of Natural History, Special Paper, Pocatello, ID.
Malde H.E., and H.A. Powers, 1962. Upper Cenozoic stratigraphy of western Snake River Plain, Idaho: Geological Society of America Bulletin, v. 73, p. 1197-1220.
Malde, H.E., H.A. Powers, and C.H. Marshall, 1963. Reconnaissance geologic map of west-central Snake River Plain, Idaho: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-373.
Matthews, S.H., 2000. Geology of Owinza Butte, Shoshone SE, and Star Lake quadrangles, Snake River Plain, southern Idaho: University of South Carolina M.S. thesis, 110 p.
O'Connor, J.E., 1993. Hydrology, hydraulics, and geomorphology of the Bonneville Flood: Geological Society of America Special Paper 274, 83 p.
Scott, W.E., 1982. Surficial geologic map of the eastern Snake River Plain and adjacent areas, 111 to 115 W., Idaho and Wyoming: U.S. Geological Survey Miscellaneous Investigations Series Map I-1372, scale 1:250,000.
Tauxe, Lisa, Casey Luskin, Peter Selkin, Phillip Gans, and Andy Calvert, 2004. Paleomagnetic results from the Snake River Plain: contribution to the time-averaged field global database: Geochemistry Geophysics Geosystems (G3), v. 5, no.8, Q08H13 DOI 10.1029/2003GC000661.



Base map scanned from USGS film positive, 1992.
Topography by photogrammetric methods from aerial photographs taken 1962. Updated from aerial photographs taken 1987 and field checked. Map edited 1992. Conflicts may exist between some updated features and previously mapped contours.
Polyconic projection, 1927 North American Datum.
10,000-foot grid ticks based on Idaho coordinate system, central zone.
1000-meter Universal Transverse Mercator grid ticks, zone 11.



Field work conducted 2003.
This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program.
Digital cartography by Jane S. Freed at the Idaho Geological Survey's Digital Mapping Lab.
Map version I-29-2005.
PDF (Acrobat Reader) map may be viewed online at www.idahogeology.org.