

PRELIMINARY GEOLOGIC MAP OF THE HAYDEN LAKE QUADRANGLE, KOOTENAI COUNTY, IDAHO

James L. Browne
2000

This Technical Report is a reproduction of independent mapping by James L. Browne of Coeur d'Alene, Idaho. Its content and format may not conform to agency's standards.



Description of Map Units

Quaternary	Qal	Recent alluvium -- clay, silt, sand, and gravel in stream valleys.
	Qgf	Sand, gravel, cobbles, and boulders transported by outburst floods from glacial Lake Missoula.
Tertiary	Tcr	Columbia River basalt -- tholeiitic basalt flows and accompanying pillow basalt and palagonitic tuff.
	Tog	Old gravel -- tan to orangish colored silt, sand, gravel, and cobbles. Gravel and cobbles are sub-angular to subrounded, both resting on and interbedded with Columbia River basalts. Grades westward into Latah Formation in places. Probably derived mainly from highlands to the east.
Tertiary or Cretaceous	Tl	Latah Formation -- clay, silt, and sand interbedded with Columbia River basalt flows. Grades eastward into old gravel (Tog) in places. Probably derived mainly from highlands to the west.
	TMsg	Stocks and other irregular bodies of granodioritic to quartz monzonitic composition.
pre-Cambrian Middle Proterozoic	Wallace Formation	
	PCW1	Laminated to thin-bedded gray to brown dolomite, laminated dolomite with black argillite and minor thin-bedded, interlayered quartzite or siltite with black argillite. Equivalent to Unit 4 in the Clark Fork, Idaho district. (Harrison and Jobn, 1963)
	PCW2	Thin-bedded to laminated gray quartzite and siltite, rusty-weathering dolomitic quartzite and feroan dolomite with prominent but quantitatively minor black argillite bedding planes. Lower part of unit consists in places of finely interlaminated black argillite, feroan dolomite and gray quartzite-siltite. Equivalent to Units 2 and 3 in the Clark Fork, Idaho district (Harrison and Jobn, 1963)
	Lowest Wallace, equivalent to Unit 1 in the Clark Fork, Idaho district (Harrison and Jobn, 1963), subdivided as follows:	
	PCW1a	Interbedded feroan dolomite, green argillite and gray to rusty-weathering (ferroan dolomite bearing) siltite-quartzite. Few thin beds of black argillite. Dolomite is dominant constituent.
	PCW1b	Thin- to medium-bedded black argillite, gray to rusty-weathering quartzite-siltite and feroan dolomite.
	PCW1c	Thin- to medium- bedded green and purple argillite, gray to rusty-weathering quartzite-siltite and feroan dolomite. Argillite and quartzite-siltite are dominant constituents.
PCsr	St. Regis Formation -- green and purple argillite with green siltite and gray to tan and greenish impure quartzite. Contains a little black argillite in places. Quartzite is especially prominent in the lower half of the formation, where it is more abundant than argillite and siltite. Few wisps and thin beds of feroan dolomite, especially toward the top. Top of unit placed at lower contact of prominent feroan dolomite layers or quartzite beds with substantial feroan dolomite content.	
PCr	Revet Formation -- thin- to thick-bedded gray to white and some greenish quartzite with thin- to medium-bedded siltite and thin-bedded greenish argillite in places. Quartzite is generally more vitreous, blocky, and less susceptible to weathering than underlying Burke Formation. Argillite increases toward formation's top, which is placed at the last medium to thick bedded white quartzite layers.	
PCb	Burke Formation -- thin- to thick-bedded gray to greenish, subvitreous siltite with abundant argillite and siltite-argillite, especially in the lower one third. Contains numerous layers of quartzite, which in a few thin beds resembles vitreous Revet quartzite, in the upper third to half of the formation. The top of the Burke is placed at the bottom of thick beds of vitreous white quartzite.	

Reference

Harrison, J.E. and Jobn, D.A., 1963. Geology of the Clark Fork Quadrangle Idaho-Montana: US Geological Survey Bulletin 1141-K. 38p.

Symbols

---	Contact, approximately located	
----	Contact, concealed	
---	Fault, approximately located	
----	Fault, concealed	
Strike and dip of beds		
$\frac{30}{\text{inclined}}$	$\frac{90}{\text{vertical}}$	$\frac{0}{\text{horizontal}}$
Strike and dip of foliation		
$\frac{40}{\text{inclined}}$	$\frac{90}{\text{vertical}}$	
\bigcirc	Area of abundant outcrop and/or roadcut exposure	
\times	Individual outcrop, roadcut exposure or diagnostic rubble	

Base map USGS digital raster graphic.
Control by USGS and NOS/NOAA.
Topography by photogrammetric methods from aerial photographs taken 1958. Field checked 1961.
Polyconic projection. 10,000-foot grid based on Idaho coordinate system, west zone.
1927 North American datum.
National geodetic datum of 1929.

UTM grid and 1961 magnetic north declination at center of sheet.

SCALE: 1:24,000
1 1/2 0 1 MILE
1 0.5 0 1 KILOMETER

Contour interval 40 feet.
Dotted lines represent 20-foot contours.

Field work 1998-1999.
Version 1-16-01.

QUADRANGLE LOCATION